

A Novel Chiral Boson

Fiorenzo Bastianelli[†]

Institute for Theoretical Physics

University of Stockholm

Vanadisvägen 9

S-113 46 Stockholm, Sweden

We introduce a new model describing a bosonic system with chiral properties. It consists of a free boson with two peculiar couplings to the background geometry which generalizes the Feigen-Fuchs-Dotsenko-Fateev construction. By choosing the two background charges of the model, it is possible to achieve any prefixed value of the left and right central charges and, in particular, obtain chiral bosonization. A supersymmetric version of the model is also given. We use the latter to identify the effective action induced by chiral superconformal matter.

[†] e-mail: fiorenzo@vana.physto.se

Bosonization is an interesting phenomenon of two dimensional field theory. It states the equality of correlation functions of given operators in the two apparently different theories of a scalar boson and a Dirac fermion [1]. In the massless case, it has been useful to extend the fermion-boson correspondence to a single chiral sector, and formulas for chiral bosonization on a general Riemann surface have been written down in [2]. However, the chiral bosonic piece has always been identified by extracting the holomorphic (in the modular parameters) square root of an effective action which corresponds to a non-chiral bosonic theory, and a direct comparison with a truly chiral bosonic theory is still absent. The reason is that it is difficult to give a lagrangian description of a chiral boson suitable for quantization. The first attempt to construct such an action was made by Siegel [3], who introduced a gauge field in order to constrain the boson to be chiral. Unfortunately, the gauge symmetry becomes anomalous upon quantization and mechanisms to cancel the anomaly have to be devised [4]. A different formulation, due to Floreanini and Jackiw [5], does not require the introduction of a gauge field, but the Lorentz invariance of the model is not manifestly realized. This is not a problem in principle, in fact one can still achieve several results like coupling the chiral boson to other systems [6], computing the gravitational anomalies [7] and showing chiral bosonization on a curved space with trivial topology (i.e. the infinite plane) [8]. However, the non-manifest Lorentz invariance (or coordinate invariance when coupled to gravity) makes it difficult to investigate the Floreanini-Jackiw model in non-trivial topological situations. In the present letter, we suggest yet a new description of a chiral boson which is different in spirit from the previous ones. The model we have in mind is that of a free scalar field coupled to the background geometry through two background charges. These couplings induce improvement terms in the energy-momentum tensor and modify the left and right central charges of the model, in a way similar to the usual boson with a background charge, discussed by Feigin-Fuchs and Dotsenko-Fateev [9]. One can choose the value of the two background charges to create a mismatch between the left and right central charges, thus obtaining a chiral system. We present only a local description of the model (more precisely, we work on the infinite plane) and adopt a Minkowskian signature for the two dimensional geometry. We conclude this letter by presenting a supersymmetric version of the chiral boson and its effective action. The latter is naturally identified with the typical effective action induced by chiral superconformal matter.

We start with the definition of the two dimensional geometry and introduce Lorentz covariant derivatives[†]

$$\nabla_a = E_a^\mu \partial_\mu + \omega_a J; \quad a = \pm \quad (1)$$

constrained by

$$[\nabla_+, \nabla_-] = RJ \quad (2)$$

The constraint (2) can be solved to get the usual spin connection ω_a as function of the vielbein field E_a^μ ($e \equiv \det e_\mu^a$, e_μ^a is the inverse of E_a^μ)

$$\omega_\pm = \mp \frac{1}{e} \partial_\mu (e E_\pm^\mu) \quad (3)$$

The scalar curvature R defined in (2) can be expressed as follows ($E_a \equiv E_a^\mu \partial_\mu$)

$$\begin{aligned} R &= R_l + R_r \\ R_l &= \nabla_+ \omega_- = (E_+ - \omega_+) \omega_- \\ R_r &= -\nabla_- \omega_+ = -(E_- + \omega_-) \omega_+ \end{aligned} \quad (4)$$

Note that R_l and R_r are not separately local Lorentz invariant. In fact, under local Lorentz (λ) and Weyl (σ) transformations defined by

$$\hat{e}_\mu^\pm = \exp(\sigma \mp \lambda) e_\mu^\pm \quad (5)$$

one obtains the following transformed quantities

$$\begin{aligned} \hat{e} \hat{R}_l &= e \left(R_l + \nabla^2 (\sigma - \lambda) \right) \\ \hat{e} \hat{R}_r &= e \left(R_r + \nabla^2 (\sigma + \lambda) \right) \end{aligned} \quad (6)$$

where $\nabla^2 \equiv \nabla_- \nabla_+ = \nabla_+ \nabla_-$ acts on scalars. Only the combination which gives the Ricci scalar $R = R_l + R_r$ is Lorentz invariant. However, we are looking for a chiral theory where gravitational anomalies are expected [10]. Moreover, it is known that one can shift possible gravitational anomalies into the Lorentz sector by using local counterterms [11]. Therefore, it is natural to use independently R_l and R_r in the construction of a model which is going to be reparametrization invariant but Lorentz (and Weyl) anomalous.

[†] J is the Lorentz generator acting on a vectors $V_\pm$ as follows $[J, V_\pm] = \pm V_\pm$; the Minkowski metric in light cone coordinates is given by $\eta_{+-} = \eta_{-+} = \frac{1}{2}$ and $\eta_{++} = \eta_{--} = 0$.

Inspired by the previous considerations, we write down the following action for a chiral boson ϕ

$$S = \frac{1}{\pi} \int d^2x e \left(\frac{1}{2} \nabla_+ \phi \nabla_- \phi + \beta_r R_r \phi + \beta_l R_l \phi \right) \quad (7)$$

To justify the claim that ϕ is a chiral boson, we begin the analysis of the model by looking at the energy-momentum. The latter is defined by $T^\mu_a = \frac{2\pi}{e} \frac{\delta S}{\delta e_\mu^a}$ and in general will not be symmetric if the theory is not local Lorentz invariant (which is the case of our model).

We obtain

$$\begin{aligned} T_{\mu a} = & e_{\mu a} \nabla_+ \phi \nabla_- \phi - e_{\mu+} \nabla_a \phi \nabla_- \phi - e_{\mu-} \nabla_+ \phi \nabla_a \phi \\ & + 2\beta_r (e_{\mu a} E_+ \nabla_- \phi - e_{\mu+} E_a \nabla_- \phi - e_{\mu-} \omega_+ \nabla_a \phi) \\ & + 2\beta_l (e_{\mu a} E_- \nabla_+ \phi - e_{\mu-} E_a \nabla_+ \phi + e_{\mu+} \omega_- \nabla_a \phi) \end{aligned} \quad (8)$$

In flat space, where $e_{\mu a} = \eta_{\mu a}$, we get the following components of the energy-momentum tensor

$$\begin{aligned} T_{++} &= -\frac{1}{2} \partial_+ \phi \partial_+ \phi - \beta_l \partial_+^2 \phi \\ T_{--} &= -\frac{1}{2} \partial_- \phi \partial_- \phi - \beta_r \partial_-^2 \phi \\ T_{+-} &= \beta_r \partial_+ \partial_- \phi = 0 \\ T_{-+} &= \beta_l \partial_- \partial_+ \phi = 0 \end{aligned} \quad (9)$$

where we have used the equation of motion evaluated in flat space, $\partial_+ \partial_- \phi = 0$, which also implies the conservation of the energy-momentum tensor. From the above expressions we see that the coupling to the background geometry has induced extra terms in the $T_{\pm\pm}$ components. Their effect is to shift the central charges of the model as follows

$$c_l = 1 + 12\beta_l^2; \quad c_r = 1 + 12\beta_r^2 \quad (10)$$

where the central charges c_l and c_r are identified from the coefficients of the anomalous terms in the operator product expansion of the energy-momentum tensor[†] [12]

$$\begin{aligned} T_{++}(x^+) T_{++}(y^+) &= \frac{c_l/2}{(x^+ - y^+)^4} + \frac{2T_{++}(y^+)}{(x^+ - y^+)^2} + \frac{\partial_+ T_{++}(y^+)}{(x^+ - y^+)} + \dots \\ T_{--}(x^-) T_{--}(y^-) &= \frac{c_r/2}{(x^- - y^-)^4} + \frac{2T_{--}(y^-)}{(x^- - y^-)^2} + \frac{\partial_- T_{--}(y^-)}{(x^- - y^-)} + \dots \end{aligned} \quad (11)$$

For $\beta_l^2 \neq \beta_r^2$ we have a chiral theory and thus a chiral boson.

[†] We use the operator product expansion $\partial_\pm \phi(x^\pm) \partial_\pm \phi(y^\pm) = \frac{1}{(x^\pm - y^\pm)^2} + \dots$

An equivalent way of describing the shift in the central charges is to compute the effective action in curved space by integrating out the bosonic field ϕ in the functional integral. This effective action can be interpreted as the generating functional of correlation functions of the energy-momentum tensor. We get

$$e^{iW[e_\mu^a]} = \int (\mathcal{D}\phi) e^{iS} \quad (12)$$

$$W[e_\mu^a] = \frac{1}{24\pi} \int d^2x e \left(R \frac{1}{\nabla^2} R + 12(\beta_l R_l + \beta_r R_r) \frac{1}{\nabla^2} (\beta_l R_l + \beta_r R_r) \right)$$

The above result is easily obtained by shifting the field $\phi \rightarrow \phi + \frac{1}{\nabla^2} (\beta_l R_l + \beta_r R_r)$ to reduce the computation to that of a standard bosonic scalar without background charges, which gives the well-known $R \frac{1}{\nabla^2} R$ effective action [13] with a coefficient of $\frac{1}{24\pi}$ in our conventions. Note that, up to *boundary terms* (we partially integrate freely assuming suitable boundary conditions at infinity) and *local terms* (e.g. $e R_l \frac{1}{\nabla^2} R_r = e \omega_- \omega_+ +$ a total derivative), one can rewrite the effective action as follows

$$W[e_\mu^a] = \frac{1}{24\pi} \int d^2x e \left((1 + 12\beta_l^2) R_l \frac{1}{\nabla^2} R_l + (1 + 12\beta_r^2) R_r \frac{1}{\nabla^2} R_r \right) \quad (13)$$

which is the effective action of a chiral system, first discussed by Leutweyler [14] for the case of a Weyl fermion (see also ref.[15] for useful formulas). To achieve central charges $c_{r,l} < 1$ one has to take imaginary couplings $\beta_{r,l}$, as in the non-chiral case [9]. In particular, the choice $\beta_l = 0$ and $\beta_r = \pm \frac{i}{2\sqrt{3}}$ reproduces the effective action of a Weyl fermion.

It is interesting to observe a relation between the chiral bosonic action (7) and the effective action induced by chiral fields (13): the action (7) is identical to what may be called the “Liouville action” for the Lorentz mode of the vielbein. To see this, parametrize the class of Lorentz-equivalent vielbeins by $e_\mu^\pm = \exp(\mp\lambda) \hat{e}_\mu^\pm$, where $\hat{e}_\mu^\pm$ is a reference vielbein. Plugging this parametrization in the action (13) one gets

$$W[e_\mu^a] = W[\hat{e}_\mu^a] + \frac{1}{24\pi} \int d^2x \hat{e} \left[(2 + \beta_l^2 + \beta_r^2) \lambda \hat{\nabla}^2 \lambda + 2(1 + \beta_r^2) \hat{R}_r \lambda - 2(1 + \beta_l^2) \hat{R}_l \lambda \right] \quad (14)$$

(the inclusion of a local counterterm of the type $\Delta S = \int d^2x e \omega_- \omega_+$ in the action (13) would bring in just some modifications of the various coefficients). Thus, we see that by adjusting the various coupling constants, the “Liouville action” for the Lorentz mode can be used to describe a chiral boson. This is similar to the usual Liouville action for the Weyl mode, which is used to represent non-chiral matter with arbitrary central charge

(note that one is typically interested in $c < 1$ and must use imaginary couplings in the Liouville action).

A supersymmetric version of the model can easily be constructed by repeating the previous steps in superspace [16]. The geometry of the latter is locally defined by introducing Lorentz covariant derivatives[†]

$$\nabla_A = E_A^M D_M + \omega_M J; \quad A = +, -, ++, -- \quad (15)$$

constrained by

$$\begin{aligned} \{\nabla_+, \nabla_+\} &= 2i\nabla_{++} \\ \{\nabla_-, \nabla_-\} &= -2i\nabla_{--} \\ \{\nabla_+, \nabla_-\} &= RJ \end{aligned} \quad (16)$$

The scalar curvature can again be expressed as a sum of chiral pieces

$$\begin{aligned} R &= R_l + R_r \\ R_l &= \nabla_+ \omega_- = \left(E_+ - \frac{1}{2}\omega_+\right)\omega_- \\ R_r &= \nabla_- \omega_+ = \left(E_- + \frac{1}{2}\omega_-\right)\omega_+ \end{aligned} \quad (17)$$

It is useful to give here the formulas for the fermionic spin connections

$$\omega_{\pm} = \mp \frac{2}{e} D_M (e E_{\pm}^M) \quad (18)$$

from which one can derive the following useful scaling properties of R_l and R_r under Weyl (Σ) and Lorentz (Λ) transformations

$$\begin{aligned} \hat{E}_{\pm}^M &= \exp(\Sigma \pm \Lambda) E_{\pm}^M \\ \hat{e} \hat{R}_l &= e \left(R_l - 2\nabla^2 (\Sigma + \Lambda) \right) \\ \hat{e} \hat{R}_r &= e \left(R_r - 2\nabla^2 (\Sigma - \Lambda) \right) \end{aligned} \quad (19)$$

where now $\nabla^2 \equiv \nabla_+ \nabla_- = -\nabla_- \nabla_+$. The superspace equivalent of the action (7) looks similar

$$S = \frac{1}{2\pi} \int d^2 x d^2 \theta e \left(\frac{1}{2} \nabla_+ \phi \nabla_- \phi + \beta_l R_l \phi + \beta_r R_r \phi \right) \quad (20)$$

[†] $D_M = \{D_+, D_-, \partial_{++}, \partial_{--}\}$ are the covariant derivatives of rigid superspace, where $D_{\pm} = \frac{\partial}{\partial \theta^{\pm}} \pm i\theta^{\pm} \frac{\partial}{\partial x^{\pm\pm}}$. The Lorentz generator now acts as follows $[J, V_{\pm}] = \pm \frac{1}{2} V_{\pm}$, $[J, V_{\pm\pm}] = +V_{\pm\pm}$.

and give rise to an affective action identical in form to (13)

$$W[e_M^A] = \frac{1}{32\pi} \int d^2x d^2\theta e \left((1 + 8\beta_l^2) R_l \frac{1}{\nabla^2} R_l + (1 + 8\beta_r^2) R_r \frac{1}{\nabla^2} R_r + a\omega_+\omega_- \right) \quad (21)$$

where the last term represent the ambiguity of adding local terms consistently with reparametrization invariance (and rigid Lorentz transformations). This effective action is obtained by shifting the field ϕ and by making use of the non-chiral effective action due to a scalar superfield [17], which is still of the type $R \frac{1}{\nabla^2} R$. It can be seen that this effective action is indeed induced by chiral supermatter. In fact, one can compute the energy-momentum tensor from the action (20) to check that it corresponds to a typical chiral superconformal system. The energy-momentum tensor is found by varying the vielbein field $\delta E_A^M \equiv H_A^B E_B^M$. One must recall that not all variation H_A^B are allowed because the constraints (16) which define the supergeometry must be preserved. There are only six independent variations which can be taken to be $H_\pm^\pm, H_\pm^{\pm\pm}, H_\pm^{\mp\mp}$. As for the remaining variations, their functional dependence is found from varying the constraints (16). The energy-momentum tensor is then identified from

$$\delta S = \frac{1}{2\pi} \int d^2x d^2\theta e (-1)^A T_A^B H_B^A \quad (22)$$

where H_B^A runs over the above set of independent variations. In flat space and using the equation of motion, we get the following non-zero components of the energy-momentum tensor

$$\begin{aligned} T_{++}^- &= -\frac{1}{2} D_{++} \phi D_+ \phi - \beta_l D_{++} D_+ \phi \\ T_{--}^+ &= \frac{1}{2} D_{--} \phi D_- \phi + \beta_r D_{--} D_- \phi \end{aligned} \quad (23)$$

It is straightforward to check that the above components of the energy-momentum tensor generate two copies of a superVirasoro algebra with central charges $\hat{c}_l = 1 + 8\beta_l^2$ and $\hat{c}_r = 1 + 8\beta_r^2$ respectively, where the central charge $\hat{c}$ is recognized from the operator product expansion of the spin $|\frac{3}{2}|$ components of the super energy-momentum tensor (see [18] for notations)

$$T(Z_1)T(Z_2) = \frac{\hat{c}/4}{z_{12}^3} + \frac{(3/2)\theta_{12}T(Z_2)}{z_{12}^2} + \frac{(1/2)D_2T(Z_2) + \theta_{12}\partial_2T(Z_2)}{z_{12}} + \dots \quad (24)$$

and it is related to the Virasoro central charge c by $c = \frac{3}{2}\hat{c}$.

To summarize, we have introduced a model of chiral bosons which reproduce the correct anomalies of chiral conformal matter in spaces with trivial topology. We hope that

the clear geometrical formulation will help to discuss the model also in topologically more interesting situations, like Riemann surfaces. The model can easily be supersymmetrized by working in superspace. The supersymmetric version has allowed us to identify the effective action induced by chiral superconformal matter, a result which we believe new by itself.

References

- [1] S. Coleman, Phys. Rev. **D11** (1975) 2088;
S. Mandelstam, Phys. Rev. **D11** (1975) 3026.
- [2] E. Verlinde and H. Verlinde, Nucl. Phys. **B288** (1987) 357;
T. Eguchi and H. Ooguri, Phys. Lett. **B187** (1987) 127;
L. Alvarez-Gaumé, J.B. Bost, G. Moore, P. Nelson and C. Vafa, Comm. Math. Phys. **112** (1987) 503.
- [3] W. Siegel, Nucl. Phys. **B238** (1984) 307.
- [4] C. Hull, Phys. Lett. **B206** (1988) 235; **B212** (1988) 347.
- [5] R. Floreanini and R. Jackiw, Phys. Rev. Lett. **59** (1987) 1973.
- [6] F. Bastianelli and P. van Nieuwenhuizen, Phys. Lett. **B217** (1989).
- [7] F. Bastianelli and P. van Nieuwenhuizen, Phys. Rev. Lett. **63** (1989) 728.
- [8] F. Bastianelli, Phys. Lett. **B254** (1991) 427.
- [9] B.L. Feigin and D.S. Fuchs, Funct. Anal. Appl. **16** (1982) 114 ;
V.I. Dotsenko and V.A. Fateev, Nucl.Phys. **B240** [FS12] (1984) 312; **B251** [FS13] (1985) 691.
- [10] L. Alvarez-Gaumé and E. Witten, Nucl. Phys. **B234** (1984) 269.
- [11] W. Bardeen and B. Zumino, Nucl. Phys. **B244** (1984) 421.
- [12] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, Nucl. Phys. **B241** (1984) 333.
- [13] A.M. Polyakov, Phys. Lett. **B103** (1981) 207.
- [14] H. Leutwyler, Phys. Lett. **B153** (1985) 65.
- [15] F. Bastianelli, P. van Nieuwenhuizen and A. Van Proeyen, Phys. Lett. **B253** (1991) 67.
- [16] P.S. Howe, J. Phys. **A12** (1979) 393.
- [17] A.M. Polyakov, Phys. Lett. **B103** (1981) 211;
E. Martinec, Phys. Rev. **D28** (1983) 2604.
- [18] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986) 93.